

## Solutions to Practice Question 2: Modeling a Non-periodic Context

2. The temperature of Shania's butternut squash soup when it is first removed from the stove ( $t = 0$ ) is  $185^\circ$  Fahrenheit. Ten minutes later ( $t = 10$ ), Shania uses a thermometer to determine that the temperature of the soup has cooled to  $130^\circ$  Fahrenheit.

The temperature of Shania's bowl of soup can be modeled by the function  $S$  given by  $S(t) = ab^t + 72$ , where  $S(t)$  is the temperature of the soup, in degrees Fahrenheit ( $^\circ\text{F}$ ), and  $t$  is the number of minutes since the soup was removed from the stove.

(A)	(i)	Since $S(0) = 185$ and $S(10) = 130$ , two equations to find $a$ and $b$ are: $185 = ab^0 + 72$ $130 = ab^{10} + 72$	1 pt
	(ii)	$185 = ab^0 + 72 \Rightarrow a = 185 - 72 = 113$ $130 = 113b^{10} + 72 \Rightarrow b = \left(\frac{130 - 72}{113}\right)^{\frac{1}{10}} = 0.935481$	1 pt
(B)	(i)	$\frac{S(10) - S(0)}{10 - 0} = \frac{130 - 185}{10} = -5.5^\circ\text{F per minute}$	1 pt
	(ii)	The secant line between $t = 0$ and $t = 10$ passes through the points $(0,185)$ and $(10,130)$ and is given by $y = 185 - 5.5t$ or $y = 130 - 5.5(t - 10)$ . For $t = 8$ , $y = 185 - 5.5(8) = 141^\circ$ . The temperature of the soup 8 minutes after it was removed from the stove was approximately $141^\circ$ .	1 pt
	(iii)	The temperature values estimated using the average rate of change are the y-coordinates of the points on the secant line between $(0,185)$ and $(10,130)$ . Since $S$ is an exponential decay function that is decreasing and concave up on its entire domain, the secant line between $t = 0$ and $t = 10$ will be above the graph of $S$ on the entire interval $(0, 10)$ . Thus, estimates for the temperature of the soup between $t = 0$ and $t = 10$ found using the average rate of change will be strictly greater than the values predicted by the model.	1 pt
(C)		The value of $M$ represents the ambient temperature of the room the soup is in and $y = M$ represents the horizontal asymptote of $S$ . Analyzing the end behavior of $S$ shows that $M = 72$ . Shania finishes the soup by the time the temperature is $10^\circ$ above the room temperature, or $82^\circ$ . Solving $S(t) = 82$ gives $t = 36.357$ . Thus, an appropriate domain would be $0 \leq t \leq 36.357$ or $0 \leq t \leq 36$ .	1 pt